

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8/18 (KRT8.18/1346), 0.2mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal BOT-BNUB1346-100

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 8/18 (KRT8.18/1346), 0.2mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1346-100 |
| Hersteller Artikelnummer | BNUB1346-100 |
| Alternativnummer | BOT-BNUB1346-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Cytoskeleton preparation from HeLa cells |
| Produktbeschreibung | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb recognizes all simple epithelia including glandular epithelium, for example thyr... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [KRT8.18/1346] |
| Molekulargewicht | 52.5 kDa (CK8), 45 kDa (CK18) |

| | |
|------------------------|--|
| UniProt | P05787 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |