

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Alpha-1-Antitrypsin (SERPINA1) (Hepatocellular & Histiocytic Marker) (AAT/1378), 0.2mg/mL, IgG1, Clone: [AAT/1378], Mouse, Monoclonal BOT-BNUB1378-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Alpha-1-Antitrypsin (SERPINA1) (Hepatocellular & Histiocytic Marker) (AAT/1378), 0.2mg/mL, IgG1, Clone: [AAT/1378], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUB1378-500  |
| Hersteller Artikelnummer | BNUB1378-500  |
| Alternativnummer         | BOT-BNUB1378-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human Alpha-1-Antitrypsin (AAT) protein fragment (exact sequence is proprietary)  |
| Produktbeschreibung      | This antibody recognizes a protein of 54 kDa, which is identified as Alpha-1 antitrypsin (AAT). AAT is secreted and is a serine protease inhibitor whose targets include elastase, plasmin, thrombin, trypsin, chymotrypsin, and plasminogen activator. De... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [AAT/1378]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 54 kDa   |
| Isotyp                 | IgG1   |
| UniProt                | <a href="#">P01009</a>   |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide  |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |