

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Factor XIIIa (Coagulation Factor XIIIa Chain) (F13A1/1448), 0.2mg/mL, Clone: [F13A1/1448], Mouse, Monoclonal BOT-BNUB1448-500

| | |
|--------------------------|---|
| Artikelname | Factor XIIIa (Coagulation Factor XIIIa Chain) (F13A1/1448), 0.2mg/mL, Clone: [F13A1/1448], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1448-500 |
| Hersteller Artikelnummer | BNUB1448-500 |
| Alternativnummer | BOT-BNUB1448-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is proprietary) |
| Produktbeschreibung | This antibody recognizes a protein of 83 kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macropha... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [F13A1/1448] |

| | |
|------------------------|---|
| Molekulargewicht | 83 kDa |
| UniProt | P00488 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |