

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Lewis Y (Tumor Marker) (LWY/1463), 0.2mg/mL, Clone: [LWY/1463], Mouse, Monoclonal BOT-BNUB1463-500

| | |
|--------------------------|---|
| Artikelname | Lewis Y (Tumor Marker) (LWY/1463), 0.2mg/mL, Clone: [LWY/1463], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1463-500 |
| Hersteller Artikelnummer | BNUB1463-500 |
| Alternativnummer | BOT-BNUB1463-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human colon carcinoma cells |
| Produktbeschreibung | This antibody recognizes a carbohydrate epitope present on tumor-associated Lewis Y antigen (Fucalpha1-2Galbeta1-4/3GlcNAcbeta). Lewis Y is expressed in large bowel tumors and colorectal carcinomas. It may be useful in the classification of human ren... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [LWY/1463] |
| Molekulargewicht | Multiple |

| | |
|------------------------|---|
| UniProt | Not Applicable |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |