

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100A4 (Marker of Tumor Metastasis) (S100A4/1481), 0.2mg/mL, Clone: [S100A4/1481], Mouse, Monoclonal BOT-BNUB1481-100

| | |
|--------------------------|---|
| Artikelname | S100A4 (Marker of Tumor Metastasis) (S100A4/1481), 0.2mg/mL, Clone: [S100A4/1481], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1481-100 |
| Hersteller Artikelnummer | BNUB1481-100 |
| Alternativnummer | BOT-BNUB1481-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human S100A4 protein |
| Produktbeschreibung | S100A4 belongs to the S100 super-family of proteins containing 2 EF-hand calcium-binding domains. S100 genes include at least 25 members, including S100A1-S100A18, trichohyalin, filaggrin, repetin, S100P, and S100Z. S100A4 exerts its function via dir... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [S100A4/1481] |
| Molekulargewicht | 10-12 kDa |

| | |
|------------------------|---|
| UniProt | P26447 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |