

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Negative Control for Rabbit Monoclonal Antibodies (NCRBM/1520R), 0.2mg/mL, Clone: [NCRBM/1520R] BOT-BNUB1520-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Negative Control for Rabbit Monoclonal Antibodies (NCRBM/1520R), 0.2mg/mL, Clone: [NCRBM/1520R]   |
| Artikelnummer            | BOT-BNUB1520-500  |
| Hersteller Artikelnummer | BNUB1520-500  |
| Alternativnummer         | BOT-BNUB1520-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Rabbit  |
| Produktbeschreibung      | To serve as a true negative control for various procedures, this rabbit monoclonal IgG should be used under the identical conditions and at the same dilution as the rabbit monoclonal antibody being used for a positive reaction. Available purified, or... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [NCRBM/1520R]   |
| Puffer                   | PBS, 0.05% BSA, 0.05% azide   |
| Quelle                   | Animal  |

|                        |  |
|------------------------|--|
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |
|------------------------|--|