

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **BTLA(BTLA/1528), 0.2mg/mL, Clone: [BTLA/1528], Mouse, Monoclonal BOT-BNUB1528-100**

|                          |   |
|--------------------------|---|
| Artikelname              | BTLA(BTLA/1528), 0.2mg/mL, Clone: [BTLA/1528], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNUB1528-100  |
| Hersteller Artikelnummer | BNUB1528-100  |
| Alternativnummer         | BOT-BNUB1528-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human BTLA protein  |
| Produktbeschreibung      | B and T lymphocyte attenuator (BTLA) is a member of the CD28 Ig superfamily of inhibitory co-receptors, along with CTLA-4 and PD-1. However, unlike CTLA-4 and PD-1, BTLA does not interact with B7-Ig family receptors. Rather, it is engaged by herpesvi... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [BTLA/1528]   |
| Molekulargewicht         | 33 kDa  |
| UniProt                  | Q7Z69   |

|                        |  |
|------------------------|--|
| Puffer                 | PBS, 0.05% BSA, 0.05% azide  |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |