

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Desmoglein-3 (Squamous Cell Marker) (DSG3/1535), 0.2mg/mL, Clone: [DSG3/1535], Mouse, Monoclonal BOT-BNUB1535-100**

Artikelname	Desmoglein-3 (Squamous Cell Marker) (DSG3/1535), 0.2mg/mL, Clone: [DSG3/1535], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1535-100
Hersteller Artikelnummer	BNUB1535-100
Alternativnummer	BOT-BNUB1535-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human DSG3 extracellular protein fragment (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of 130 kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein c...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[DSG3/1535]

Molekulargewicht	130 kDa
UniProt	<a href="#">P32926</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p>