

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/1611), 0.2mg/mL, IgG1, Clone: [LPFS2/1611], Mouse, Monoclonal BOT-BNUB1611-100

Artikelname	CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/1611), 0.2mg/mL, IgG1, Clone: [LPFS2/1611], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1611-100
Hersteller Artikelnummer	BNUB1611-100
Alternativnummer	BOT-BNUB1611-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human CD22 protein (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[LPFS2/1611]
Molekulargewicht	120 kDa

Isotyp	IgG1
UniProt	P26842
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user