

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **S100A (Melanoma Marker) (S1/61), 0.2mg/mL, Clone: [S1/61], Mouse, Monoclonal BOT-BNUB1695-100**

Artikelname	S100A (Melanoma Marker) (S1/61), 0.2mg/mL, Clone: [S1/61], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1695-100
Hersteller Artikelnummer	BNUB1695-100
Alternativnummer	BOT-BNUB1695-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Human
Immunogen	Purified human S-100 protein conjugated to methylated BSA
Produktbeschreibung	S100A and S100B proteins are two members of the S100 family of calcium binding proteins. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epitope located on the alpha-ch...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[S1/61]
Molekulargewicht	10-12 kDa

UniProt	<a href="#">P23297</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user