

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 15 (KRT15) (Basal Cell Marker) (KRT15/1699), 0.2mg/mL, Clone: [KRT15/1699], Mouse, Monoclonal BOT-BNUB1699-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 15 (KRT15) (Basal Cell Marker) (KRT15/1699), 0.2mg/mL, Clone: [KRT15/1699], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUB1699-500  |
| Hersteller Artikelnummer | BNUB1699-500  |
| Alternativnummer         | BOT-BNUB1699-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Bovine, Human, Rat  |
| Immunogen                | Recombinant human KRT15 protein   |
| Produktbeschreibung      | Keratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I keratin (keratins 9 to 23) and one type II keratin (keratins 1 to 8). Keratins demonstrate tissue and differentiation spec... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [KRT15/1699]  |
| Molekulargewicht         | 52 kDa  |

|                        |   |
|------------------------|---|
| UniProt                | <a href="#">P19012</a>  |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (frozen or formalin) 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |