

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) (IG1707R), 0.2mg/mL, Clone: [IG1707R], Rabbit, Monoclonal BOT-BNUB1707-100

| | |
|--------------------------|---|
| Artikelname | Human IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) (IG1707R), 0.2mg/mL, Clone: [IG1707R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUB1707-100 |
| Hersteller Artikelnummer | BNUB1707-100 |
| Alternativnummer | BOT-BNUB1707-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Purified human Ig Gamma Chain |
| Produktbeschreibung | This antibody recognizes a protein of 75 kDa, identified as gamma heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [IG1707R] |
| Molekulargewicht | 75 kDa |

| | |
|------------------------|---|
| UniProt | P01861 , P01860 , P01859 , P01857 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |