

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD61 / Integrin beta-3 / Platelet Glycoprotein IIIa (Platelet Marker) (ITGB3/1713), 0.2mg/mL, Clone: [ITGB3/1713], Mouse, Monoclonal BOT-BNUB1713-500**

|                          |   |
|--------------------------|---|
| Artikelname              | CD61 / Integrin beta-3 / Platelet Glycoprotein IIIa (Platelet Marker) (ITGB3/1713), 0.2mg/mL, Clone: [ITGB3/1713], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNUB1713-500  |
| Hersteller Artikelnummer | BNUB1713-500  |
| Alternativnummer         | BOT-BNUB1713-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human ITGB3 protein   |
| Produktbeschreibung      | This antibody reacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the alpha-V-chain (CD51) to form vitronectin receptor, or with the alpha-IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The C... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [ITGB3/1713]  |
| Molekulargewicht         | 105 kDa & 90 kDa  |

|                        |  |
|------------------------|--|
| UniProt                | <a href="#">P05106</a>   |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide  |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Immunohistology (frozen or formalin) 1-2 ug/mL Optimal dilution for a specific application should be determined by user |