

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (F8/86), 0.2mg/mL, IgG1, Clone: [F8/86], Mouse, Monoclonal BOT-BNUB1716-100

Artikelname	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (F8/86), 0.2mg/mL, IgG1, Clone: [F8/86], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1716-100
Hersteller Artikelnummer	BNUB1716-100
Alternativnummer	BOT-BNUB1716-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Factor isolated from human plasma von Willebr
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[F8/86]
Molekulargewicht	250 kDa

Isotyp	IgG1
UniProt	P04275
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>