

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glycophorin A / CD235a (Erythrocyte Marker) (JC159), 0.2mg/mL, Clone: [JC159], Mouse, Monoclonal BOT-BNUB1737-500

Artikelname	Glycophorin A / CD235a (Erythrocyte Marker) (JC159), 0.2mg/mL, Clone: [JC159], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1737-500
Hersteller Artikelnummer	BNUB1737-500
Alternativnummer	BOT-BNUB1737-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Formalin fixed membrane from a Hairy cell leukemia
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 39 kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[JC159]
Molekulargewicht	39 kDa

UniProt	P02724
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user