

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Connexin 32 (Gap Junction Protein) (GJB1/1753), 0.2mg/mL, Clone: [GJB1/1753], Mouse, Monoclonal BOT-BNUB1753-500

| | |
|--------------------------|---|
| Artikelname | Connexin 32 (Gap Junction Protein) (GJB1/1753), 0.2mg/mL, Clone: [GJB1/1753], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1753-500 |
| Hersteller Artikelnummer | BNUB1753-500 |
| Alternativnummer | BOT-BNUB1753-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Recombinant human GJB1 protein |
| Produktbeschreibung | This Ab recognizes a protein of 27-32 kDa, identified as Connexin 32. The connexin family of proteins forms hexameric complexes called connexons that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin prote... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [GJB1/1753] |
| Molekulargewicht | 27-32 kDa |

| | |
|------------------------|---|
| UniProt | P08034 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Immunofluorescence 1-2 ug/mL Flow cytometry 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |