

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Bcl-X (Apoptosis Marker)(BCL2L1/1762), 0.2mg/mL, Clone: [BCL2L1/1762], Mouse, Monoclonal BOT-BNUB1762-100

Artikelname	Bcl-X (Apoptosis Marker)(BCL2L1/1762), 0.2mg/mL, Clone: [BCL2L1/1762], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1762-100
Hersteller Artikelnummer	BNUB1762-100
Alternativnummer	BOT-BNUB1762-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Recombinant human BCL2L1 protein fragment (around aa 1-135) (exact sequence is proprietary)
Produktbeschreibung	Bcl-x belongs to the Bcl-2 family of proteins and plays a crucial role in apoptosis. Bcl-X has two isoforms: Bcl-XL (long), a 26 kDa protein which suppresses cell death, and Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain whic...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[BCL2L1/1762]
Molekulargewicht	27 kDa

UniProt	Q07817
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry