

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), 0.2mg/mL, Clone: [VWF/1767], Mouse, Monoclonal BOT-BNUB1767-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)<br>(VWF/1767), 0.2mg/mL, Clone: [VWF/1767], Mouse, Monoclonal                                                                                                                             |
| Artikelnummer            | BOT-BNUB1767-500                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB1767-500                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB1767-500-500UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant fragment of human vWF protein (aa1815-1939) (exact sequence is proprietary)                                                                                                                                                                       |
| Produktbeschreibung      | von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [VWF/1767]                                                                                                                                                                                                                                                    |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 250 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| UniProt                | <a href="#">P04275</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |