

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) (PCRP-CDX2-1A3), 0.2mg/mL, IgG1, Clone: [PCRP-CDX2-1A3], Mouse, Monoclonal BOT-BNUB1920-500

| | |
|--------------------------|---|
| Artikelname | CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) (PCRP-CDX2-1A3), 0.2mg/mL, IgG1, Clone: [PCRP-CDX2-1A3], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1920-500 |
| Hersteller Artikelnummer | BNUB1920-500 |
| Alternativnummer | BOT-BNUB1920-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CDX2protein |
| Produktbeschreibung | The specificity of this monoclonal antibody to its intended target was validated by HuProt(TM)Array, containing more than 19,000, full-length human proteins.The intestine-specific transcription factors CDX1 and CDX2 are important for directing intest... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [PCRP-CDX2-1A3] |
| Molekulargewicht | 40 kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| UniProt | Q99626 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Immunoprecipitation 2-4 ug/mg protein Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |