

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Prostate Specific Acid Phosphatase (PSAP) (rACPP/1338), 0.2mg/mL, Clone: [rACPP/1338], Mouse, Monoclonal BOT-BNUB2074-100

| | |
|--------------------------|---|
| Artikelname | Prostate Specific Acid Phosphatase (PSAP) (rACPP/1338), 0.2mg/mL, Clone: [rACPP/1338], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB2074-100 |
| Hersteller Artikelnummer | BNUB2074-100 |
| Alternativnummer | BOT-BNUB2074-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human ACPP protein |
| Produktbeschreibung | This antibody recognizes a protein of 52 kDa, identified as prostate specific acid phosphatase (PSAP). This enzyme catalyzes the conversion of orthophosphoric monoester to alcohol and orthophosphate. It is synthesized under androgen regulation and is... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [rACPP/1338] |
| Molekulargewicht | 52 kDa |

| | |
|------------------------|---|
| UniProt | P15309 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |