

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD52 (Epididymis-Specific Protein 5) (CD52/2276R), 0.2mg/mL, Clone: [CD52/2276R], Rabbit, Monoclonal BOT-BNUB2276-500

| | |
|--------------------------|--|
| Artikelname | CD52 (Epididymis-Specific Protein 5) (CD52/2276R), 0.2mg/mL, Clone: [CD52/2276R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUB2276-500 |
| Hersteller Artikelnummer | BNUB2276-500 |
| Alternativnummer | BOT-BNUB2276-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Monkey |
| Immunogen | Recombinant full-length human CD52 protein |
| Produktbeschreibung | Human CD52 is expressed at high density by lymphocytes, monocytes, eosinophils, thymocytes and macrophages. It is expressed by most lymphoid derived malignancies, although expression on myeloma cells is variable. Primary antibodies are available purif... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [CD52/2276R] |

| | |
|------------------------|---|
| Molekulargewicht | 6 kDa |
| UniProt | P31358 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |