

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC18 / CD146 / MCAM (Melanoma Cell Adhesion Molecule) (rMUC18/1130), 0.2mg/mL, IgG1, Clone: [rMUC18/1130], Mouse, Monoclonal BOT-BNUB2302-100

| | |
|--------------------------|--|
| Artikelname | MUC18 / CD146 / MCAM (Melanoma Cell Adhesion Molecule) (rMUC18/1130), 0.2mg/mL, IgG1, Clone: [rMUC18/1130], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB2302-100 |
| Hersteller Artikelnummer | BNUB2302-100 |
| Alternativnummer | BOT-BNUB2302-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human MUC18 protein |
| Produktbeschreibung | Mel-CAM belongs to the immunoglobulin superfamily and functions as a Ca ²⁺ -independent cell adhesion molecule. Mel-CAM expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not norm... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [rMUC18/1130] |

| | |
|------------------------|--|
| Molekulargewicht | 130 kDa |
| Isotyp | IgG1 |
| UniProt | P43121 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA pH 8.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |