

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) (rKRT19/799), 0.2mg/mL, IgG1, Clone: [rKRT19/799], Mouse, Monoclonal BOT-BNUB2304-100**

Artikelname	Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) (rKRT19/799), 0.2mg/mL, IgG1, Clone: [rKRT19/799], Mouse, Monoclonal
Artikelnummer	BOT-BNUB2304-100
Hersteller Artikelnummer	BNUB2304-100
Alternativnummer	BOT-BNUB2304-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human KRT19 protein
Produktbeschreibung	This MAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40 kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[rKRT19/799]
Molekulargewicht	40 kDa

Isotyp	IgG1
UniProt	<a href="#">P08727</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat or ferret Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user