

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TRAcP (Tartrate-Resistant Acid Phosphatase) (Hairy Cell Leukemia Marker) (ACP5/2336R), 0.2mg/mL, Clone: [ACP5/2336R], Rabbit, Monoclonal BOT-BNUB2336-100**

|                          |   |
|--------------------------|---|
| Artikelname              | TRAcP (Tartrate-Resistant Acid Phosphatase) (Hairy Cell Leukemia Marker) (ACP5/2336R), 0.2mg/mL, Clone: [ACP5/2336R], Rabbit, Monoclonal  |
| Artikelnummer            | BOT-BNUB2336-100  |
| Hersteller Artikelnummer | BNUB2336-100  |
| Alternativnummer         | BOT-BNUB2336-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human, Mouse, Rat   |
| Immunogen                | Recombinant full-length human ACP5 protein  |
| Produktbeschreibung      | This antibody recognizes a protein of 35 kDa, which is identified as tartrate-resistant acid phosphatase (TRAcP). It exists as two isoforms (5a and 5b). This MAb reacts with both the isoforms. Serum TRAcP 5a is secreted by macrophages and dendritic c... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [ACP5/2336R]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 35 kDa  |
| UniProt                | <a href="#">P13686</a>  |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |