

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker) (TYRP1/2340R), 0.2mg/mL, Clone: [TYRP1/2340R], Rabbit, Monoclonal BOT-BNUB2340-100**

Artikelname	Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker) (TYRP1/2340R), 0.2mg/mL, Clone: [TYRP1/2340R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2340-100
Hersteller Artikelnummer	BNUB2340-100
Alternativnummer	BOT-BNUB2340-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human TYRP1 protein (around aa 257-377) (exact sequence is proprietary)
Produktbeschreibung	This antibody reacts with a 75 kDa melanocyte-specific gene product, identified as Tyrosinase-Related Protein-1 (TRP-1). It is involved in melanin synthesis. TRP1 is present on the melanosomal membranes of melanoma, normal melanocytes and nevi. Recen...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[TYRP1/2340R]

Molekulargewicht	75 kDa
UniProt	<a href="#">P17643</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user