

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD40 / TNFRSF5 / CD40L-Receptor (C40/2383), 0.2mg/mL, Clone: [C40/2383], Mouse, Monoclonal BOT-BNUB2383-100

| | |
|--------------------------|---|
| Artikelname | CD40 / TNFRSF5 / CD40L-Receptor (C40/2383), 0.2mg/mL, Clone: [C40/2383], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB2383-100 |
| Hersteller Artikelnummer | BNUB2383-100 |
| Alternativnummer | BOT-BNUB2383-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human CD40 protein (around aa 29-107) (exact sequence is proprietary) |
| Produktbeschreibung | CD40 is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [C40/2383] |

| | |
|------------------------|---|
| Molekulargewicht | 43 kDa |
| UniProt | P25942 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |