

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2427), 0.2mg/mL, Clone: [TP63/2427], Mouse, Monoclonal BOT-BNUB2427-100

Artikelname	p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2427), 0.2mg/mL, Clone: [TP63/2427], Mouse, Monoclonal
Artikelnummer	BOT-BNUB2427-100
Hersteller Artikelnummer	BNUB2427-100
Alternativnummer	BOT-BNUB2427-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human p63 protein
Produktbeschreibung	p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[TP63/2427]
Molekulargewicht	63 kDa

UniProt	Q9H3D4
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user