

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Prostate Specific Antigen (PSA) (KLK3/2871R), 0.2mg/mL, Clone: [KLK3/2871R], Rabbit, Monoclonal BOT-BNUB2871-500

| | |
|--------------------------|---|
| Artikelname | Prostate Specific Antigen (PSA) (KLK3/2871R), 0.2mg/mL, Clone: [KLK3/2871R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUB2871-500 |
| Hersteller Artikelnummer | BNUB2871-500 |
| Alternativnummer | BOT-BNUB2871-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KLK3 protein |
| Produktbeschreibung | This antibody recognizes a single protein of 33-34 kDa, identified as the prostate specific antigen (PSA). This MAb is highly specific to PSA and stains prostatic secretory and ductal epithelium in both normal and neoplastic tissues. PSA is a chymotr... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [KLK3/2871R] |
| Molekulargewicht | 33-34 kDa |

| | |
|------------------------|--|
| UniProt | P07288 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |