

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD81 / TAPA-1 (C81/2885R), 0.2mg/mL, Clone: [C81/2885R], Rabbit, Monoclonal BOT-BNUB2885-100**

Artikelname	CD81 / TAPA-1 (C81/2885R), 0.2mg/mL, Clone: [C81/2885R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2885-100
Hersteller Artikelnummer	BNUB2885-100
Alternativnummer	BOT-BNUB2885-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	Exosome Staining, FC, IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	B-Cell line derived from a Burkitt lymphoma
Produktbeschreibung	This antibody recognizes a protein of 26 kDa, identified as CD81 (Workshop VI, Code CD81.1). CD81 has a very broad cellular distribution, being expressed on T- and B-lymphocytes, NK cells, thymocytes, eosinophils, fibroblasts, epithelial and endothel...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[C81/2885R]
Molekulargewicht	26 kDa

UniProt	<a href="#">P60033</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user