

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD68 (Macrophage Marker) (C68/2908R), 0.2mg/mL, Clone: [C68/2908R], Rabbit, Monoclonal BOT-BNUB2908-500**

|                          |   |
|--------------------------|---|
| Artikelname              | CD68 (Macrophage Marker) (C68/2908R), 0.2mg/mL, Clone: [C68/2908R], Rabbit, Monoclonal  |
| Artikelnummer            | BOT-BNUB2908-500  |
| Hersteller Artikelnummer | BNUB2908-500  |
| Alternativnummer         | BOT-BNUB2908-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human LAMP4 protein   |
| Produktbeschreibung      | This antibody recognizes a glycoprotein of 110 kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [C68/2908R]   |
| Molekulargewicht         | ~110 kDa  |

|                        |   |
|------------------------|---|
| UniProt                | <a href="#">P34810</a>  |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |