

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TLE1 (Synovial Sarcoma Marker) (TLE1/2946R), 0.2mg/mL, Clone: [TLE1/2946R], Rabbit, Monoclonal BOT-BNUB2946-100

| | |
|--------------------------|---|
| Artikelname | TLE1 (Synovial Sarcoma Marker) (TLE1/2946R), 0.2mg/mL, Clone: [TLE1/2946R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUB2946-100 |
| Hersteller Artikelnummer | BNUB2946-100 |
| Alternativnummer | BOT-BNUB2946-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human TLE1 fragment (aa 175-338) (exact sequence is proprietary) |
| Produktbeschreibung | Key players in the Notch pathway are the TLE genes, which are human homologs of the Drosophila groucho gene. Groucho is a transcriptional repressor that plays a key role in neurogenesis, segmentation and sex determination. Transducin-like enhancer pr... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [TLE1/2946R] |

| | |
|------------------------|---|
| Molekulargewicht | 83 kDa |
| UniProt | Q04724 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |