

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC2 (Mucin 2) (MLP/2970R), 0.2mg/mL, Clone: [MLP/2970R], Rabbit, Monoclonal BOT-BNUB2970-100

| | |
|--------------------------|---|
| Artikelname | MUC2 (Mucin 2) (MLP/2970R), 0.2mg/mL, Clone: [MLP/2970R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUB2970-100 |
| Hersteller Artikelnummer | BNUB2970-100 |
| Alternativnummer | BOT-BNUB2970-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human MUC2 protein |
| Produktbeschreibung | This antibody recognizes a single glycoprotein of 520 kDa, identified as mucin 2 (MUC2). This MAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC3. Its epitope has been defined as GTQTP (GlyThrGlnThrPro). Mucins are high m... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [MLP/2970R] |
| Molekulargewicht | 520 kDa |

| | |
|------------------------|---|
| UniProt | Q02817 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris-HCl buffer pH 10.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |