

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tenascin C (Stromal Marker For Epithelial Malignancy) (TNC/2981R), 0.2mg/mL, Clone: [TNC/2981R], Rabbit, Monoclonal BOT-BNUB2981-500

Artikelname	Tenascin C (Stromal Marker For Epithelial Malignancy) (TNC/2981R), 0.2mg/mL, Clone: [TNC/2981R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2981-500
Hersteller Artikelnummer	BNUB2981-500
Alternativnummer	BOT-BNUB2981-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human TNC protein
Produktbeschreibung	In Western blotting, this antibody reacts with two bands of ~MW of 210 kDa and 300 kDa, identified as two isoforms of Tenascin C. Specificity of this MAb is validated by sequential immunoprecipitation with a PAb against Tenascin C. Tenascin C is a mu...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[TNC/2981R]

Molekulargewicht	210 kDa and 300 kDa
UniProt	P24821
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user