

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 0.2mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal BOT-BNUB2988-100

Artikelname	BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 0.2mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2988-100
Hersteller Artikelnummer	BNUB2988-100
Alternativnummer	BOT-BNUB2988-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human BCL10 protein fragment (around aa122-168) (exact sequence is proprietary)
Produktbeschreibung	B-cell lymphoma/leukemia 10 (BCL10) is a signaling protein that promotes apoptosis, pro-caspase-9 maturation, and activation of NF-kappa-B via NIK and IKK. BCL10 contains an N-terminal caspase recruitment domain (CARD), which is found in a number of ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[BL10/2988R]

Molekulargewicht	33 kDa
UniProt	O95999
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user