

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### CD28(C28/75), 1mg/mL, Clone: [C28/75], Mouse, Monoclonal BOT-BNUM0075-50

Artikelname	CD28(C28/75), 1mg/mL, Clone: [C28/75], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0075-50
Hersteller Artikelnummer	BNUM0075-50
Alternativnummer	BOT-BNUM0075-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC
Spezies Reaktivität	Human
Immunogen	Recombinant human CD28 protein
Produktbeschreibung	Recognizes a glycoprotein of 44-88 kDa, which is identified as CD28. It is the critical T-cell co-stimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[C28/75]
Molekulargewicht	44 kDa (monomer), 88 kDa (dimer)
UniProt	<a href="#">P10747</a>

Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user