

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MHC II (HLA-DP) (beta chain)(BRA-FB6), 1mg/mL, Clone: [BRA-FB6], Mouse, Monoclonal BOT-BNUM0199-50

Artikelname	MHC II (HLA-DP) (beta chain)(BRA-FB6), 1mg/mL, Clone: [BRA-FB6], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0199-50
Hersteller Artikelnummer	BNUM0199-50
Alternativnummer	BOT-BNUM0199-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, FC, Functional Studies, IP
Spezies Reaktivität	Human
Immunogen	Non-T, non-B human acute lymphoblastic leukemia REH6 cell line
Produktbeschreibung	Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35 kDa) and beta (27-30 kDa) chains. It reportedly reacts with B- & non-T, non-B cell lines but not w...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BRA-FB6]
Molekulargewicht	33-35 kDa

UniProt	P04440
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user