

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD6(C6/372), 1mg/mL, IgG1, Clone: [C6/372], Mouse, Monoclonal BOT-BNUM0372-50

| | |
|--------------------------|---|
| Artikelname | CD6(C6/372), 1mg/mL, IgG1, Clone: [C6/372], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0372-50 |
| Hersteller Artikelnummer | BNUM0372-50 |
| Alternativnummer | BOT-BNUM0372-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human recombinant CD6 protein |
| Produktbeschreibung | CD6 is a type I transmembrane glycoprotein that contains a 24-amino acid signal sequence, three extracellular scavenger receptor cysteine-rich (SRCR) domains, a membrane-spanning domain and a 44-amino acid cytoplasmic domain. The CD6 glycoprotein is ... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [C6/372] |
| Molekulargewicht | 90-130 kDa |
| Isotyp | IgG1 |

| | |
|------------------------|--|
| UniProt | P30203 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |