

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p40(Rabbit PAb), 1mg/mL, Clone: [PAb], Monoclonal BOT-BNUM0375-50

| | |
|--------------------------|---|
| Artikelname | p40(Rabbit PAb), 1mg/mL, Clone: [PAb], Monoclonal |
| Artikelnummer | BOT-BNUM0375-50 |
| Hersteller Artikelnummer | BNUM0375-50 |
| Alternativnummer | BOT-BNUM0375-50-50UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Bovine, Human, Mouse, Rat |
| Immunogen | A synthetic peptide (ENNAQTQFSEPQY) corresponding to aa5-17 of human p40 |
| Produktbeschreibung | p40 (p63 delta) is a marker determined to be highly specific for squamous basal cells by immunohistochemistry (IHC). The current more routinely recommended marker, p63, appears to have less specificity compared to p40, especially on squamous cell tum... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [PAb] |
| Molekulargewicht | 40 kDa |

| | |
|------------------------|--|
| UniProt | Q9H3D4 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |