

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker)(MOC-31), 1mg/mL, Clone: [MOC-31], Mouse, Monoclonal BOT-BNUM0380-50

Artikelname	Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker)(MOC-31), 1mg/mL, Clone: [MOC-31], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0380-50
Hersteller Artikelnummer	BNUM0380-50
Alternativnummer	BOT-BNUM0380-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Neuraminidase treated GLS-1 human small cell lung carcinoma cells
Produktbeschreibung	The epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cel...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MOC-31]
Molekulargewicht	40-43 kDa

UniProt	P16422
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Does not react with rat Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user