

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Cytokeratin 19(Ks19.1), 1mg/mL, Clone: [Ks19.1], Mouse, Monoclonal BOT-BNUM0394-50

Artikelname	Cytokeratin 19(Ks19.1), 1mg/mL, Clone: [Ks19.1], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0394-50
Hersteller Artikelnummer	BNUM0394-50
Alternativnummer	BOT-BNUM0394-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Human breast cancer MCF-7 cells
Produktbeschreibung	This MAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40 kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[Ks19.1]
Molekulargewicht	40 kDa
UniProt	<a href="#">P08727</a>

Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user