

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgM Immunoglobulin(Cocktail), 1mg/mL, Clone: [Cocktail], Mouse, Monoclonal BOT-BNUM0762-50

Artikelname	IgM Immunoglobulin(Cocktail), 1mg/mL, Clone: [Cocktail], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0762-50
Hersteller Artikelnummer	BNUM0762-50
Alternativnummer	BOT-BNUM0762-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant heavy chain of human IgM (IM260 & IM373), Heavy chain of human IgM (ICO-30)
Produktbeschreibung	Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[Cocktail]

Molekulargewicht	50-75 kDa
UniProt	P01871
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user