

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD20 / MS4A1 (B-Cell Marker)(L26), 1mg/mL, Clone: [L26], Mouse, Monoclonal BOT-BNUM0766-50

Artikelname	CD20 / MS4A1 (B-Cell Marker)(L26), 1mg/mL, Clone: [L26], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0766-50
Hersteller Artikelnummer	BNUM0766-50
Alternativnummer	BOT-BNUM0766-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Human tonsil B cells
Produktbeschreibung	This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[L26]
Molekulargewicht	33-37 kDa

UniProt	P11836
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user