

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 17(KRT17/778), 1mg/mL, Clone: [KRT17/778], Mouse, Monoclonal BOT-BNUM0778-50**

Artikelname	Cytokeratin 17(KRT17/778), 1mg/mL, Clone: [KRT17/778], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0778-50
Hersteller Artikelnummer	BNUM0778-50
Alternativnummer	BOT-BNUM0778-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Bovine, Goat, Human, Porcine, Rat
Immunogen	Recombinant full-length human KRT17 protein
Produktbeschreibung	Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[KRT17/778]
Molekulargewicht	46 kDa

UniProt	<a href="#">Q04695</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user