

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 18(KRT18/835), 1mg/mL, IgG1, Clone: [KRT18/835], Mouse, Monoclonal BOT-BNUM0835-50

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 18(KRT18/835), 1mg/mL, IgG1, Clone: [KRT18/835], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0835-50 |
| Hersteller Artikelnummer | BNUM0835-50 |
| Alternativnummer | BOT-BNUM0835-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KRT18 protein |
| Produktbeschreibung | This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [KRT18/835] |
| Molekulargewicht | 45 kDa |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| UniProt | P05783 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Does not react with mouse, rat, sheep, hamster, cow, dog or pig, others not known Optimal dilution for a specific application should be determined by user</p> |