

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p34 / cdk1(POH-1), 1mg/mL, Clone: [POH-1], Mouse, Monoclonal BOT-BNUM0853-50

Artikelname	p34 / cdk1(POH-1), 1mg/mL, Clone: [POH-1], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0853-50
Hersteller Artikelnummer	BNUM0853-50
Alternativnummer	BOT-BNUM0853-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Mink, Monkey
Immunogen	Purified recombinant human p34cdc2 fusion protein
Produktbeschreibung	Recognizes a 34 kDa protein, which is identified as cyclin dependent kinase 1 (cdk1) or p34cdc2 protein kinase. cdk1 / p34cdc2 plays a crucial role during cell division and is most active during mitosis. It is predominantly localized in the nucleus. ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[POH-1]
Molekulargewicht	34 kDa
UniProt	P06493

Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse, rat, Xenopus, Drosophila, S. pombe, or S. cerevisiae, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>