

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TTF-1 / NKX2.1(8G7G3/1 + NX2.1/690), 1mg/mL, Clone: [8G7G3/1 NX2.1/690], Mouse, Monoclonal BOT-BNUM0907-50

| | |
|--------------------------|---|
| Artikelname | TTF-1 / NKX2.1(8G7G3/1 + NX2.1/690), 1mg/mL, Clone: [8G7G3/1 NX2.1/690], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0907-50 |
| Hersteller Artikelnummer | BNUM0907-50 |
| Alternativnummer | BOT-BNUM0907-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Rat full length TTF-1 recombinant protein (8G7G3/1), Recombinant TTF-1 protein (NX2.1/690) |
| Produktbeschreibung | Recognizes a protein of 40 kDa, identified as Thyroid transcription factor-1 (TTF-1). TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver,... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [8G7G3/1 NX2.1/690] |

| | |
|------------------------|--|
| Molekulargewicht | 40 kDa |
| UniProt | P43699 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |