

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Tyrosinase(T311 + OCA1/812), 1mg/mL, Clone: [T311 OCA1/812], Mouse, Monoclonal BOT-BNUM0908-50**

Artikelname	Tyrosinase(T311 + OCA1/812), 1mg/mL, Clone: [T311 OCA1/812], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0908-50
Hersteller Artikelnummer	BNUM0908-50
Alternativnummer	BOT-BNUM0908-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant tyrosinase protein (T311), Recombinant human TYR protein (OCA1/812)
Produktbeschreibung	Recognizes a cluster of proteins between 70-80 kDa, identified as tyrosinase. Occasionally a minor band at 55 kDa is also detected. This MAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-cont...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[T311 OCA1/812]

Molekulargewicht	70-80 kDa
UniProt	<a href="#">P14679</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, or 10 mM citrate buffer, pH6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user