

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(IIIIE2.34), 1mg/mL, Clone: [IIIIE2.34], Mouse, Monoclonal BOT-BNUM0934-50

Artikelname	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(IIIIE2.34), 1mg/mL, Clone: [IIIIE2.34], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0934-50
Hersteller Artikelnummer	BNUM0934-50
Alternativnummer	BOT-BNUM0934-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human vWF fragment spanning aa 845-949
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[IIIIE2.34]
Molekulargewicht	250 kDa

UniProt	P04275
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Western blot: 0.5-1.0 ug/mL Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user