

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A63-C/A9), 1mg/mL, Clone: [A63-C/A9], Mouse, Monoclonal BOT-BNUM0938-50

| | |
|--------------------------|---|
| Artikelname | Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A63-C/A9), 1mg/mL, Clone: [A63-C/A9], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0938-50 |
| Hersteller Artikelnummer | BNUM0938-50 |
| Alternativnummer | BOT-BNUM0938-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Neuraminidase-treated human red blood cells |
| Produktbeschreibung | This antibody recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivit... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [A63-C/A9] |
| Molekulargewicht | Multiple |

| | |
|------------------------|---|
| UniProt | Not Applicable |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at 37°C Immunohistochemistry: frozen: 0.5-1.0 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |