

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myeloid-Associated Differentiation Marker(MYADM/971), 1mg/mL, IgG1, Clone: [MYADM/971], Mouse, Monoclonal BOT-BNUM0971-50

Artikelname	Myeloid-Associated Differentiation Marker(MYADM/971), 1mg/mL, IgG1, Clone: [MYADM/971], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0971-50
Hersteller Artikelnummer	BNUM0971-50
Alternativnummer	BOT-BNUM0971-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Primate
Immunogen	Recombinant human MYADM protein
Produktbeschreibung	Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellul...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MYADM/971]
Molekulargewicht	Unknown

Isotyp	IgG1
UniProt	Q96S97
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user